

DO YOU KNOW ?

1. The most common malignancy seen in ataxia telangiectasia patients
 - a. breast cancer
 - b. carcinoma of tongue
 - c. lymphomas
 - d. melanoma
 2. A "fingertip unit" of ointment (from distal crease to tip of index finger) is approximately
 - a. 0.2 to 0.3 gm.
 - b. 0.3 to 0.4 gm
 - c. 0.4 to 0.5 gm
 - d. 0.5 to 0.6 gm
 3. Wood's light is ultraviolet light of wavelength
 - a. 365 nm
 - b. 340 to 360 nm
 - c. 320 to 340nm
 - d. 320 nm
 4. The most prominent clinical feature of Bloom's syndrome
 - a. Characteristic facies
 - b. high-pitched voice
 - c. photosensitivity
 - d. small body size
 5. The risk of contracting HIV through percutaneous injury is estimated at
 - a. 0.2% per exposure
 - b. 0.3%
 - c. 0.4%
 - d. 0.5%
 6. The features of seronegative spondyloarthropathies include
 - a. genetic susceptibility
 - b. immune complexes and complement consumption
 - c. involvement of periarticular structures or enthesopathy
 - d. variable involvement of axial skeleton
 7. Cutaneous atrophy due to topical corticosteroid results in alterations in the
 - a. epidermis
 - b. dermis
 - c. vascular elements of skin
 - d. subcutaneous tissue
 8. The first-generation retinoids include
 - a. acitretin
 - b. etretinate
 - c. isotretinoin
 - d. tretinoin
 9. The strength of trichloroacetic acid used for superficial chemical peeling.
 - a. 1-10%
 - b. 10-25%
 - c. 25-35%
 - d. 35-50%
 10. The pathognomonic skin markers for Cowden syndrome
 - a. multiple facial trichilemmomas
 - b. multiple palmoplantar punctate lesions
 - c. multiple oral 'cobblestone' lesions
 - d. multiple sclerotic fibromas
- M.P Binitha**

1.c 2.c 3.a 4.d 5.b 6.a,c,d 7.a,b,c 8.c,d 9.b 10.a,d

ANSWERS