

**LETTERS TO THE EDITOR****Kyrlé's disease and vitamin D resistant rickets in chronic renal failure****To the Editor:**

Kyrlé's disease is one of the perforating dermatoses and a rare chronic disorder of unknown aetiology. The pathogenic event in this condition is a disorder of epidermal keratinization. Clinically Kyrlé's disease is characterized by hyperkeratotic para-follicular or follicular papules, with central cone shaped plug which can be removed easily, usually involving the extensor surface of extremities. Histopathology reveals a heavy keratotic, partly parakeratotic plug containing basophilic debris lying in an invagination of the epidermis.<sup>1</sup> Several authors have reported the association of Kyrlé's disease with diabetes mellitus, renal failure, hyperlipidaemia, liver diseases, congestive cardiac failure with an infective process and with abnormal vitamin A metabolism.<sup>2</sup>

A 16-year-old male patient presented with asymptomatic hyperkeratotic parafollicular and follicular discrete papules with cone shaped removable central plug since four months. The lesions first appeared on the extensor aspect of thighs and upper extremities and gradually involved the buttocks, back and lower abdomen. The patient had abnormal presence of posterior urethral valves leading to bilateral hydronephrosis and chronic renal failure since the age of seven years. His blood urea and serum creatinine were raised throughout in all repeated investigations. Later on he developed vitamin D re-

sistant rickets. His serum calcium was 7.26 mg% and serum phosphorus was 5.3 mg%. There was no family history of similar illness. Haemogram, blood sugar, liver function test, serum lipid profile, X-ray chest and ECG were within normal limits. X-rays of skeletal system showed characteristic changes of rickets. Histopathology of skin lesions was suggestive of Kyrlé's disease.

So in this patient of chronic renal failure there was a simultaneous association of vitamin D resistant rickets and Kyrlé's disease.

Samir Shah

JN Dave

NS Vora

Benny Cardoso

Rajesh Goel

From the Department of Skin and V.D, Bapu Nagar General Hospital (ESIS), Ahmedabad-380024, India

**Clinical and bacteriological study of pyoderma in Jodhpur-Western Rajasthan****To the Editor:**

Pyogenic skin infection is one of the common conditions. The universal use of antibiotics has produced changes in bacterial flora of man and increased resistance by the micro-organism.<sup>1</sup> In a study conducted in our department on 200 patients with purulent skin infection, various predisposing factors along with organism causing pyoderma and their sensitivity to commonly used antibiotics were considered.

Diagnosis was made on clinical grounds and those patients who had not received any antibiotics in the past 15 days were taken in this study. Laboratory investigations included smear for Gram's staining, bacteriological cul-