

DO YOU KNOW

1. The mosaic artefact or pattern of irregular polygons seen in potassium hydroxide preparation for examination of fungus is due to-
 - a) lipid drops between cells
 - b) bubbles in the specimen
 - c) fibres on the slide
 - d) KOH crystals
2. Among monozygotic twins, if one twin has psoriasis, the chance of the other being affected is-
 - a) 63%
 - b) 73%
 - c) 83%
 - d) 93%
3. The drugs most likely to cause angioedema-
 - a) angiotensin-converting enzyme (ACE) inhibitors
 - b) beta adrenergic blockers
 - c) calcium channel blockers
 - d) diuretics
4. Histologically, the infiltrate of lichen nitidus is-
 - a) fibrinoid
 - b) focal
 - c) granulomatous
 - d) lichenoid
5. Drugs that can cause linear IgA bullous dermatosis-
 - a) barbiturates
 - b) captopril
 - c) lithium
 - d) vancomycin
6. An acute, superficial folliculitis composed mainly of eosinophils is seen in the biopsy specimen of-
 - a) candidiasis of infancy
 - b) erythema toxicum neonatorum
 - c) infantile acropustulosis
 - d) transient neonatal pustular melanosis
7. Diseases that may be distributed in the "seborrheic areas" include-
 - a) acne vulgaris
 - b) Darier's disease
 - c) pemphigus foliaceus
 - d) psoriasis
8. Direct immunofluorescence findings in porphyria cutanea tarda-
 - a) homogenous IgG at dermo-epidermal junction and around blood vessels
 - b) linear IgG and IgA at DEJ
 - c) linear C3, IgG and IgA at DEJ
 - d) IgG and C3 in intercellular spaces of epidermis
9. The drugs commonly causing drug-induced LE-
 - a) bleomycin
 - b) hydralazine
 - c) procainamide
 - d) warfarin
10. Conditions associated with leukocytoclastic vasculitis-
 - a) erythema ab igne
 - b) erythema elevatum diutinum
 - c) Henoch - Schonlein purpura
 - d) Kawasaki disease

9. b, c 10. b, c

1. a 2. b 3. a 4. c 5. b, c, d 6. b 7. b, c, d 8. a

Answers

MP Binitha