

Immunohistochemistry of COVID-19 skin




Figure 1: Painting of immunohistochemistry in skin of COVID-19 patients. TRPA1 channels are expressed in red, while Mas-1 channels are expressed in green

This is an acrylic painting inspired by the immunohistochemistry stains of the epidermis and dermis of patients with COVID-19 (Fig. 1). COVID-19 comes with a variety of dermatological conditions, including some that are pruritic. The immunohistochemistry stains depicted here stain for transient receptor potential cation channel subfamily A member 1 (TRPA1) and Mas-1-related G-protein coupled receptor, receptors believed to be involved in the transmission of itch.

Financial support and sponsorship
Nil.

Conflict of interest
There are no conflicts of interest.

Teresa Ju

Dr. Phillip Frost Department of Dermatology and Cutaneous Surgery,
University of Miami, Miami, Florida, United States.

Corresponding author:

Ms. Teresa Ju,
Dr. Phillip Frost Department of Dermatology and
Cutaneous Surgery, University of Miami,
Miami, Florida, United States.
ju.teresa17@gmail.com

How to cite this article: Ju T. Immunohistochemistry of COVID-19 skin. Indian J Dermatol Venereol Leprol 2023;89:332.

Received: March, 2022 **Accepted:** June, 2022 **Epub Ahead of Print:** October, 2022 **Published:** March, 2023

DOI: 10.25259/IJDVL_326_2022 **PMID:** ***

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, transform, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.