

## Constricting band following incomplete condom removal



**Figure 1: Cleavage at coronal sulcus**

A 28-year-old male had progressive penile ulceration for 50 days. After protected intercourse with his wife, he noted pain followed by ulceration with difficulty during urination. On examination, a deep cleft separated the glans from the shaft of penis. A constricting band was found embedded at the most distal part of the cleavage [Figure. 1]. On attempting to remove the band, it was revealed to be the proximal rolled part of a condom which had burrowed into the coronal sulcus.

**Joginder Kumar, V. Ramesh**

Department of Dermatology and STD,  
VM Medical College and Safdarjang Hospital,  
New Delhi, India

**Address for correspondence:** Dr. Joginder Kumar,  
Department of Dermatology and STD,  
VM Medical College and Safdarjang Hospital,  
New Delhi, India.  
E-mail: jogikataria@gmail.com

### Access this article online

Quick Response Code:



Website:

[www.ijdv.com](http://www.ijdv.com)

DOI:

10.4103/0378-6323.157452

**How to cite this article:** Kumar J, Ramesh V. Constricting band following incomplete condom removal. Indian J Dermatol Venereol Leprol 2015;81:542.

**Received:** January, 2015. **Accepted:** February, 2015. **Source of Support:** Nil. **Conflict of Interest:** None declared.