CASE NOTES

"TUBERCULOUS ULCER GLANS PENIS-A CASE NOTE" With Lupus Vulgaris

Ву

RAMAPEOSAD CHOSH, M.B.B.s., Research Worker AJIT K. DUTTA, M.B.B.S., D. V. (Madras) Clinical Tutor SOURIN GHOSH, MBBs, LRCP (Lond) FRCS, (Eng.), Professor.

Department of Venereo ogy and Sexual Disorders, Medical College Hospitals, Calcutta-12

Since our case note on Tuberculous ulcer glans penis (Ind. J. Derm. Vener. Oct-Dec. 1962) another case with lupus vulgaris reported for treatment.

JA, 30, MM, Worker in a cotton mills, married 8 years, 3 living children teetotaler came with a history of sore penis (4 months) associated with eruptions lower lip (6 weeks).

On examination: as above.

Histological Section:—Biopsy from Penile ulcer confirmed our diagnosis of Tubercular origin.

Chest X-Ray: P. A. view-"Infiltration Rt. upper zone".

Urine:—No abnormality – culture – negative for Acid fast Bacilli STS (V. D. R. L.) – Reactive 1:32

Biopsy:—Donovan bodies not found.

E. S. R. (Westergren)—1st hour-10 m.m. 2nd hour 24 m.m. mean average 11 m.m.

Pyelography.—(I. V.) - no abnormality detected.

Skin biopsy from lower lip.—" Structure suggestive of Lupus vulgaris".

The patient was given a routine course of Streptodihydrostreptomycin and INAH combination. Sore and "Lupus" healed. At present he is being followed up by Chest Department and V. D. Out-door.