

Pearly penile papules



Figure 1: Multiple whitish pink tiny papules in a few rows with central dotted vessels

A 20-year-old man with no significant medical history presented with an 8-year history of multiple papules on the glans of his circumcised penis and had been diagnosed as condyloma acuminata.

Clinical examination showed multiple 1–2 mm, smooth, dome-shaped, skin-colored papules in a single or double row along the corona of the glans penis. No ulcers or inguinal lymphadenopathy was noted. Dermoscopy demonstrated whitish pink tiny papules in a few rows with central dotted vessels [Figure 1]. A diagnosis of pearly

Access this article online	
Quick Response Code:	Website: www.ijdvl.com
	DOI: 10.4103/0378-6323.190869

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Hwang S, Kim GM. Pearly penile papules. Indian J Dermatol Venereol Leprol 2017;83:125-6.

Received: April, 2016. **Accepted:** June, 2016.

penile papules was made. We explained the benign nature of these lesions to the patient and no treatment was required.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

Sewon Hwang, Gyong Moon Kim

Department of Dermatology, St. Vincent's Hospital, College of Medicine,
The Catholic University of Korea, Seoul, Republic of Korea

Address for correspondence: Dr. Gyong Moon Kim,
Department of Dermatology, St. Vincent's Hospital, College of Medicine,
The Catholic University of Korea, Jungbu-daero 93, Paldal-gu,
Suwon-si, 442-723, Seoul, Republic of Korea.
E-mail: gyongmoonkim@catholic.ac.kr