

Named cells in dermatology

Fiona F. Sequeira, Suneil Gandhi¹, Usha Kini², Ishwar Bhat

INTRODUCTION

This article briefly describes the named cells in dermatology. It's purpose is to serve as a ready reckoner to the postgraduates during their quiz preparations. They include the following:

A. General category

1. Normal cutaneous anatomy: Langerhans cells, Glomus cells, Veil cells, Mast cells
2. Bullous disorders: Tzanck cells
3. Eczematous disorders: Tadpole cells
4. Metabolic and storage disorders: Gargoyle cells, Gaucher's cell, Xanthoma cells
5. Tumors: Basalioma cells, Basophilic and shadow cells, Doughnut cells, Flower cells, Granular cells, Halo cells, Hibernoma cells, Paget cells, Sezary cells, Signet ring cells, Spider web cells, Strap cells, Reed Sternberg cells, Vulvar Clear Cells of Toker, Floret giant cells
6. Histiocytic disorders: Touton giant cells

Table 1: Under general and infectious category

Name	Description/location/size	LM/EM	Stains/markers	Observed in/significance
Langerhans cell	Named after Paul Langerhans. Is a pale-staining non-keratinocyte, with a convoluted nucleus. Its cytoplasm contains vimentin intermediate filaments and small rod- or racket-shaped structures called Langerhans cell granules or Birbeck granules ^[1]	LM EM	H and E CD 1a S-100 Langerin (CD 207)	Are distributed in the epidermis (preference for a suprabasal position), oral cavity, esophagus and vagina. ^[1] They serve as the antigen-processing and presenting cells. ^[1] They are reduced in the epidermis of patients with diseases like psoriasis, sarcoidosis and contact dermatitis, and increased in langerhans cell histiocytosis.
Glomus cell	They are large polyhedral cells with a round, centrally located nucleus and a pale eosinophilic cytoplasm. ^[2] The cells are of smooth muscle origin. They comprise of specialized vascular structures called glomus formations. ^[2] These formations have an arterial and venous segment without intervening capillaries	LM EM	H and E Myosin Vimentin	Glomus tumors and glomangiomas arise from the arterial segment of the cutaneous glomus, the Suquet-Hoyer canal. ^[2] Glomus formations are special arteriovenous shunts that appear to play a role in thermoregulation. ^[2]
Veil cell	These are flat adventitial cells with distinctive nuclear structure and prominent cytoplasmic processes that surround the dermal microvessels. ^[3] They lie external to the wall, demarcating the vessel from the surrounding dermis. They express factor XIIIa and Von Willebrand factor ^[3]	LM Phase contrast EM	H and E	It is strategically positioned in the organization of the microvessel to participate in the control of bleeding at the microvascular level. ^[4]

(Continued)

Departments of Dermatology and ²Pathology, St John's Medical College and Hospital, Bangalore, Karnataka, ⁴Department of Dermatology, JN Medical College, Belgaum, India

Address for correspondence: Dr. Fiona F. Sequeira, Department of Dermatology, St John's Medical College and Hospital, Koramangala, Bangalore, Karnataka - 560 034, India. E-mail: dr_fiona@rediffmail.com

Access this article online

Quick Response Code:



Website:

www.ijdvl.com

DOI:

10.4103/0378-6323.93650

How to cite this article: Sequeira FF, Gandhi S, Kini U, Bhat I. Named cells in dermatology. Indian J Dermatol Venereol Leprol 2012;78:207-16.

Received: August, 2011. **Accepted:** November, 2011. **Source of Support:** Nil. **Conflict of Interest:** None declared.

Table 1: (Continued)

Name	Description/location/size	LM/EM	Stains/markers	Observed in/significance
Mast cell	Are round to spindle in shape with a unilobed round to oval centrally positioned nucleus. Its size varies up to 5–13 μm . They contain in their cytoplasm granules that range from 0.3 to 0.8 μm ^[5] Mast cells synthesize an impressive repertoire of mediators, some of them including histamine, heparin, tryptase, chymase, carboxypeptidase, neutrophil chemotactic factor and eosinophilic chemotactic factor of anaphylaxis ^[5]	LM EM	H and E, Metachromatic staining with toluidine blue, methylene blue, alcian blue and Giemsa: Red-purple, Leder stain, Mast cell tryptase and Chymase CD117	In the skin, mast cells are present in greatest density in the papillary dermis, near the DEJ, in sheaths of epidermal appendages, and around blood vessels and nerves of the subpapillary plexus. They are also common in the subcutaneous fat ^[1] Activated mast cells are the primary effector cells in the onset of an allergic reaction. They play a role in the defense against parasites, stimulate chemotaxis, activation, and proliferation of eosinophils, promote phagocytosis, alter vasotension and vascular permeability and stimulate connective tissue repair and angiogenesis ^[1] Their numbers are increased in mastocytosis.
Tastzellen cells, Merkel cells	Named after the German anatomist Friedrich Sigmund Merkel. These non-keratinocytes are oval in shape, have a pale-staining cytoplasm containing bundles of intermediary filaments and osmiophilic granules. The nucleus is large and lobulated, and the margins of these cells project cytoplasmic “spines” toward the keratinocytes ^[1]	LM EM	Silver impregnation, CAM 5.2 or AE1/AE3, CK 20, Neuron-specific enolase (NSE)	They are found both in hairy skin and in the glabrous skin of the digits, lips, regions of the oral cavity and the outer root sheath of the hair follicle. Merkel cells are slow-adapting, type I mechanoreceptors. Merkel cell carcinoma is a malignant tumor of Merkel cells ^[6] .
Tzanck cells/ acantholytic cells/mourning edged cells	It is named after Arnault Tzanck, a French dermatologist. These are large, round keratinocytes with an enlarged hyperchromatic nucleus, hazy or absent nucleoli and abundant basophilic cytoplasm. The basophilic staining is deeper peripherally on the cell membrane (“mourning edged” cells) due to the cytoplasm’s tendency to get condensed at the periphery, leading to a perinuclear halo. ^[7] In contrast to pemphigus vulgaris, the acantholytic cells in pemphigus foliaceus and pemphigus erythematosus often have a hyalinized cytoplasm that corresponds to the dyskeratosis seen in tissue sections	LM	H and E, Giemsa, Papanicolaou stain, Wright, Methylene blue, Toluidine blue	Pemphigus. This is a form of primary acantholysis. The other examples include Darier’s disease, transient acantholytic dermatoses and warty dyskeratoma. When acantholysis occurs secondary to some alteration/damage to epidermal cells, the term secondary acantholysis is used. E.g., bullous impetigo, herpes simplex/zoster and squamous cell carcinoma.
Tadpole cells	Are cells with a “tadpole-like” shape. Presence of more than 10 tadpole cells in a Tzanck smear at $\times 100$ magnification is a diagnostic indicator for spongiotic vesicular dermatitis ^[8,9]	LM	H and E	Spongiotic vesicular dermatitis. E.g., allergic contact dermatitis.
Gargoyle cells/ Hurler cells	A large oval or polygonal cell, 20 μm in diameter, with a pale central nucleus, the cytoplasm appearing clear on H and E staining. When stained with toluidine blue, metachromatic granules are seen in the cytoplasm. ^[10] On EM, the cytoplasm of the clear cells is seen to be filled with large irregular clear vacuoles varying in size from 0.5 μm to 2 μm . The membranes of the vacuoles are often broken and the ruptured ends curl. Occasionally, mitochondria and cisterns of endoplasmic reticulum are interspersed between the vacuoles	LM EM	H and E, Toluidine blue, Alcian blue	Are seen in Hurler syndrome.

(Continued)

Table 1: (Continued)

Name	Description/location/size	LM/EM	Stains/markers	Observed in/significance
Gaucher cells	The disease is named after the French doctor Philippe Gaucher. Gaucher cells are glucocerebroside-laden macrophages measuring 50–60 µm in diameter. On electron microscopy, they contain tubular cytoplasmic inclusions that are considered to be transformed secondary lysosomes. The fibrillar ceramide accumulations are extensive enough to give the cytoplasm a distinctive "crinkled blue tissue paper" appearance ^[11,12]	LM EM	Wright stain, PAS+, Iron stains	Gaucher's disease is an inherited disorder of ceramide metabolism in which an enzyme deficiency leads to the accumulation of glucocerebroside in macrophages of the bone marrow, reticuloendothelial system but not the skin. These cells are not diagnostic of Gaucher disease. Cutaneous manifestation of this condition include patchy pigmentation of the skin and petechiae.
Xanthoma cells	Often within the same lesion it's appearance can vary from eosinophilic to clear to foamy or to multinucleated giant cells, which are often referred to as Touton giant cells ^[13]	LM EM	H and E, Fat stains such as oil red O or Sudan black B are required to visualize lipid droplets on frozen sections	Various xanthomatoses, familial hyperlipoproteinemia types IIa, III and V, cerebrotendinous xanthomatosis, Wolman's disease, Tangier disease, Hand-Schuller-Christian disease and normolipidemic cutaneous xanthomatosis.
Basalioma cells	Trichoepitheliomas contain an epithelial component composed of uniform small basophilic cells looking like epidermal basal cells with round nuclei and cytoplasm. They are arranged in strands and lobules. They display peripheral palisading and sometimes contain horn cysts or calcifications ^[14,15]	LM	H and E	Trichoepitheliomas, BCC.
Basophilic and shadow or ghost cells	Pilomatricomas are sharply demarcated tumors that possess two types of cells: Basophilic cells and eosinophilic shadow cells. The former have round or elongated, deeply basophilic hyperchromatic nuclei and scanty cytoplasm so that nuclei lie close to each other. The latter are anucleate, eosinophilic, keratinized cells having a distinct border and possessing a central unstained area as a shadow of the lost nucleus. These cells frequently calcify ^[16]	LM	H and E	Pilomatricomas.
Doughnut cells	They are the intranuclear cytoplasmic inclusions in the large anaplastic mononuclear cells. ^[17] These inclusions are the result of deep cytoplasmic invaginations into the nuclei. Nuclear indentation >70% of the total nuclear diameter is the criterion to term it as a half-doughnut cell ^[18]	LM	H and E Papanicolaou, May-Grunwald, Giemsa, CD30+, Clusterin	Anaplastic large cell lymphoma.
Flower cells or clover leaf cells	Are peripheral blood atypical lymphocytes characterized by having irregular nuclear contours, prominent nucleoli, condensed chromatin and deeply basophilic, occasionally vacuolated cytoplasm ^[19]	LM	H and E, CD25+	Adult T cell lymphoma (ATL) is an aggressive mature T-cell malignancy caused by HTLV-I. It affects the skin in 43–72% of the cases. The disease manifests itself in the skin as erythroderma, plaques, papules, nodules and tumors. ^[20] These so-called flower cells are considered pathognomonic of ATL and are seen in the peripheral blood of these patients. ^[21]
Granular cells	Are large polygonal cells in the dermis with distinct cell membranes and prominent eosinophilic granular cytoplasm. The nuclei are usually small, round to oval and centrally located ^[22]	LM	H and E, Granules are PAS+ diastase-resistant	Granular cell tumor, considered to be of peripheral nerve sheath origin can involve various parts of the body (skin and tongue being the most common organs involved).

(Continued)

Table 1: (Continued)

Name	Description/location/size	LM/EM	Stains/markers	Observed in/significance
Halo cells	Intraepidermal lymphocytes that are larger than the normal lymphocytes with convoluted, cerebriform nuclei and a clear rim of cytoplasm ^[23]	LM	H and E	Are seen in mycosis fungoides.
Hibernoma-like cells, Mulberry cells, multivesicular cells	Are large (average- 54 microns) fat cells containing a central nucleus with a multivacuolated scanty granular, eosinophilic cytoplasm. ^[24] The main histopathologic differentiation of adult adipose tissue from hibernoma cells is the presence of a clear cytoplasm and an eccentric nucleus [Figure 1]	LM	H and E	An uncommon neoplasm derived from brown fat, ^[25] are seen mostly in adults, with a predilection for the thigh, back, axilla, neck and intrathoracic region.
Paget cells	Are round or oval cells that appear larger than the keratinocytes. They have a vacuolated amphophilic, cytoplasm with a hypertrophic nucleus, scanty nuclear chromatin and a large nucleolus. No desmosomal attachments are seen between Paget cells and the adjacent keratinocytes ^[26]	LM	H and E, PAS+	Found in Paget's disease of the breast.
Sezary cells, Lutzner cells, Mycosis cell, Cerebriform lymphocytes, Atypia cellulues monstreuses or Monster cell	First described by Albert Sézary. Are atypical lymphocytes with a grooved or cerebriform nucleus. The nuclear chromatin is condensed in a patchy distribution beneath the nuclear membrane. The nucleoli may be prominent. The cytoplasm is sparse and appears as a narrow rim around the nucleus. Mitochondria (often appearing clumped), rough endoplasmic reticulum, polysomes and cytoplasmic fibrils are seen. The cytoplasmic fibrils appear serpentine on electron microscopic examination. The smaller versions of these cells are called Lutzner cells and the larger ones as classical Sezary cells ^[27,28]	LM EM	Wright, Giemsa, PAS	Seen in mycosis fungoides and Sezary syndrome (both in tissue and in blood).
Signet ring cells	These cells have a single large vacuole full of mucin. This vacuole displaces the hyperchromatic nucleus to the periphery. ^[29] The pool of mucin in a signet ring cell mimics the appearance of a finger hole, and the nucleus mimics the appearance of the face of the ring in profile [Figure 2]	LM	H and E, Alcian blue, Mucicarmin	Are seen in mucin-producing adenocarcinomas.
Spider web cells	Are polygonal or ovoid tumor cells that exhibit cytoplasmic vacuoles due to their glycogen content. Residual strands of cytoplasm between the vacuoles define spider web cells ^[30,31]	LM	H and E	Adult rhabdomyomas.
Strap cells	Cells with elongated eosinophilic cytoplasm ^[32]	LM	H and E	Are suggestive but not diagnostic of rhabdomyosarcoma.
Reed Sternberg cells	They are named after Dorothy Reed Mendenhall and Carl Sternberg. Are large cells that are either multinucleated or have a bilobed nucleus (resembling an "owl's eye" appearance) with prominent eosinophilic inclusion-like nucleoli ^[33,34] [Figure 3]	LM	H and E, CD30, CD15	The presence of these cells is pathognomonic for the diagnosis of classical Hodgkin's lymphoma (HL). Other than HL, cells resembling Reed-Sternberg cells may be present in other B and T cell lymphomas, carcinomas, melanomas and sarcomas. The prevalence of skin involvement in HL is approximately 0.5–3.4%.

(Continued)

Table 1: (Continued)

Name	Description/location/size	LM/EM	Stains/markers	Observed in/significance
Vulvar clear cells of Tokier, intraepidermal glandular cells	Clear cells of Tokier are intraepithelial cells with clear to pale staining cytoplasm and bland cytologic features. They are found in the epidermis of the nipple, immediately adjacent to the openings of the lactiferous ducts, along the basal layer of the epidermis or scattered in the stratum Malpighii. Tokier cells are distributed as single units or as small aggregates that occasionally form small lumina. They have been considered a primitive or germinative cell of the lactiferous duct or interepithelial extension of the lactiferous duct cells ^[35,36]	LM	H and E, CK ₇ , EMA	Found in approximately 10% of normal nipples. Clear cells of Tokier have generated interest as a possible precursor cell of primary intraepidermal extramammary Paget's disease, as morphologically they appear to be a benign counterpart of the malignant cells of Paget's disease.
Floret giant cells	These are multinucleated giant cells that have marginal, often overlapping nuclei and an eosinophilic center ^[37]	LM	H and E, Vimentin, CD34	Seen in pleomorphic lipoma, giant cell collagenoma, giant cell fibroblastoma, giant cell angiofibroma, neurofibromatosis type 1.
Touton giant cells	These cells have a peripheral wreath-like arrangement of nuclei, with amphophilic cytoplasm centrally and pale, foamy cytoplasm at the periphery of the cell outside the nuclei ^[38]	LM	H and E	Seen in xanthoma, disseminated xanthoma, juvenile xanthogranuloma, necrobiotic xanthogranuloma and dermatofibroma
Hobnail cells	Are plump endothelial cells with scanty cytoplasm and enlarged prominent, hyperchromatic nuclei with one or two minute nucleoli. These endothelial cells are disposed in a monolayer, with nuclei bulging into the vascular lumina, resulting in a characteristic hobnail or matchstick appearance ^[39]	LM	H and E	Hobnail hemangioma is a rare benign vascular tumor. It usually appears as a single, small, well-circumscribed, angiomatous or brownish, flat or exophytic skin lesion.
LE cell	The LE cell is the end result of either phagocytosis of distorted nuclear material by a polymorphonuclear leukocyte or autolysis of one or more lobes of the nucleus ^[40,41] The LE cell contains within its cytoplasm the LE body, which appears as a homogenous, pale blue to deep-purplish material pushing the nucleus of the phagocyte to one side of the cell [Figure 4].	LM	Romanowsky stains	SLE. LE cells are not usually found in peripheral blood, although the LE cell phenomenon could form in the buffy coat of peripheral blood after a period of incubation.
Tart cells	A tart cell is a monocyte that has engulfed an intact lymphocyte nucleus. It differs from an LE cell because the secondary nucleus retains its definite chromatin structure ^[40,42]	LM	Romanowsky stains	It is not disease specific.
Podophyllin cells	Bizarre, enlarged epithelial cells with either a clumped pyknotic nucleus or with the nuclear chromatin disintegrated and dispersed as small granules throughout the cytoplasm of the cell. These "podophyllin cells" resemble carcinoma cells, but may be distinguished from malignancy by the absence of cytologic atypia and multinucleate giant cells ^[43]	LM	H and E	Seen after the application of podophyllin to normal skin, verruca vulgaris and condylooma accuminata.
Mantle cells/ corps ronds	Corps ronds are dykeratotic keratinocytes with a small pyknotic nucleus, acidophilic cytoplasm and a perinuclear halo ("mantle cells") ^[7] . They are seen in the stratum granulosum and upper malpighian layer	LM	H and E	Darier disease.
Half and half cells	These are transformed keratinocytes which show increased synthetic activity in one half of the cell, and fibrillar degeneration in the other half of the cell (resembling that of colloid bodies). ^[44]	EM	Phosphotungstic acid	These cells are seen in lichen planus, and are believed to provide a useful clue to the early changes occurring in colloid body formation.

(Continued)

Table 1: (Continued)

Name	Description/location/size	LM/EM	Stains/markers	Observed in/significance
Bean bag cells	They are macrophages that become engorged by phagocytosis of erythrocytes, lymphocytes and cell fragments ^[45]	LM	H and E	Seen in histiocytic cytophagic panniculitis.
Sunburn cells	Are pink, dead or dying keratinocytes characteristic of mammalian epidermis after exposure to UVC and UVB radiation or UVA radiation in the presence of psoralens. These cells have a pyknotic nucleus and a shrunken glassy eosinophilic cytoplasm and are seen scattered throughout the epidermis ^[46]	LM	H and E	Seen in sunburns and PUVA-induced acute phototoxicity.
Balloon cells	These are polyhedral cells with ample microvesicular cytoplasm, (ultrastructurally corresponding to enlarged vacuolated melanosomes) sparse to absent pigment, distinct cytoplasmic membranes and small, hyperchromatic, round to slightly scalloped central nuclei and rare intranuclear vacuoles or pseudoinclusions ^[47,48]	LM	H and E	Balloon cell nevus.
Pekin cells, Reiter cells	Are macrophages 40 µm in size. They contain one or more intact polymorphonuclear leucocytes and deep staining basophilic globular structures within the cytoplasm ^[49]	LM	H and E	In Reiter's disease (50–70% of synovial fluid specimens) as well as any other inflammatory synovial disease.
Clue cells	Are squamous epithelial cells with a large number of coccobacillary organisms, <i>Gardnerella vaginalis</i> and other anaerobic bacteria densely attached in clusters to their surfaces, giving them a granular appearance. The cytoplasm appears fuzzy (like shading with black pencil) and the edges of the squamous epithelial cells, which normally have a sharply defined cell border, become indistinct or stippled ^[50] [Figure 5]	LM	Papanicolaou, Gram	Seen in bacterial vaginosis. To be significant for bacterial vaginosis (BV), more than 20% of the epithelial cells on the wet mount should be clue cells.
Downey cells/ Turk cells	Named after Hal Downey. Are atypical lymphocytes, which, unlike leukemic cells, show extreme variability of size and shape. They have abundant pale blue cytoplasm. The nucleus may be irregular or indented with lightly stained chromatin. Cytoplasmic edges may be indented by other blood cells, leading to a scalloped appearance called Dutch skirting ^[51-53]	Oil immersion	H and E, Giemsa	Seen in the peripheral blood of infectious mononucleosis patients. The detection of at least 10% atypical lymphocytes on a peripheral blood smear in a patient with mononucleosis has a sensitivity of 75% and a specificity of 92% for the diagnosis of infectious mononucleosis.
Greenblatt and Pund cells	Are large mononuclear cells measuring 25–90 µm in diameter. The nucleus of the cell is eccentrically placed, vesicular or pyknotic. There are <20 intracytoplasmic vacuoles containing Donovan bodies in either young non-capsulated or mature capsulated forms ^[54,55]	LM	Giemsa, Leishman, Wright	Seen in granuloma inguinale.
Lepra cells/ Virchow cells	Named after Rudolph Virchow. Lepra cells when studied by means of vital staining are found to be derived from the histiocyte of the infected tissue. Their protoplasm, which forms the greater part of the cell, is filled with vacuoles, and contains enormous numbers of lepra bacilli. The nucleus is usually single and is, as a rule, pressed to one side by the vacuoles and bacilli that crowd the cell body. In some instances, these vacuoles are so numerous that there is little actual protoplasm remaining, the cell body having the appearance of a foam. ^[56] In time and with antimycobacterial therapy, degenerate bacilli accumulate in the macrophages, the so-called lepra cells, which then have a foamy or vacuolated cytoplasm [Figure 6]	LM	Modified Fite Faraco stain	They are observed in lepromatous leprosy They resemble xanthoma cells and, on staining with fat stains, are shown to contain lipid, largely neutral fat, and phospholipids rather than cholesterol ^[57]

(Continued)

Table 1: (Continued)

Name	Description/location/size	LM/EM	Stains/markers	Observed in/significance
Mickulicz cells	A large histiocyte measuring 10–100 μm in diameter. It has a pale, vacuolated cytoplasm. Within the cytoplasm of these cells, one finds many <i>Klebsiella bacilli</i> ^[58]	LM	Giemsa, Gram, Warthin-starry silver, PAS+	The characteristic cell in rhinoscleroma.
Von Hansemann cells	Named after David Paul Von Hansemann. Are large macrophages with abundant foamy eosinophilic cytoplasm and a prominent eccentric, hyperchromatic round nucleus. Variable numbers of intracytoplasmic, concentrically laminated, round–ovoid, basophilic inclusions are appreciated and are referred to as Michaelis–Gutmann bodies ^[59]	LM	H and E, PAS-positive, Diastase resistant	Seen in malakoplakia.
Langhans giant cells	Named after Theodor Langhans, a German pathologist. They are formed by the fusion of epithelioid cells and contain multiple nuclei arranged in a horseshoe-shaped pattern in the cell periphery or are arranged circumferentially ^[60]	LM	H and E	Seen in tuberculoid granulomas. E.g., tuberculoid leprosy, cutaneous tuberculosis, gumma of syphilis and sporotrichosis.
Wright cells	Are large macrophages that contain numerous protozoa grouped together in a typical "swarm of bees" fashion within the weakly basophilic cytoplasm ^[61]	LM	May Grunwald, Giemsa	Seen in cutaneous and mucosal leishmaniasis.
Warthin Finkeldey cells	A type of giant cell that vary in size, but can contain as many as 100 nuclei per cell along with cytoplasmic and nuclear inclusions ^[62]	LM	H and E	Pathognomic for measles infection. Found in reticuloendothelial system, exanthem, Koplik spots and respiratory secretions.
Koilocytes	Are human papilloma virus-infected vacuolated keratinocytes that bear a basophilic pyknotic nucleus surrounded by a clear halo and pale-staining cytoplasm. They are located in the upper stratum malpighii and in the granular layer. They contain few or no keratohyaline granules ^[63]	LM	H and E	Verruca vulgaris.
Mitosoid cells	Are keratinocytes with irregularly shaped hyperchromatic nuclei that simulate mitoses. They are in fact degenerated nuclei without mitotic ability ^[64]	LM	H and E	In focal epithelial hyperplasia (Heck disease).

LM: Light microscopy, EM: Electron microscopy




Figure 1: Hibernoma. Large polyhedral fat cells containing eosinophilic granular cytoplasm (x200)




Figure 2: Signet ring cells in an adenocarcinoma. Sheet of large cells with vacuolated cytoplasm pushing the single hyperchromatic nucleus to the periphery (x400)




Figure 3: Reed Sternberg cells in a case of Hodgkin's lymphoma. Note the bilobed nucleus with a resemblance to an "owl's eye" with prominent eosinophilic inclusion-like nucleoli (x200)




Figure 4: LE cells. Distorted nuclear material in the cytoplasm of a polymorphonuclear leukocyte causing the nucleus to be compressed and pushed to the periphery (x1000)




Figure 5: Clue cells. Squamous epithelial cells with a large number of coccobacillary organisms, *Gardnerella vaginalis* and other anaerobic bacteria densely attached in clusters to their surfaces give them a granular appearance (x400)




Figure 6: Lepra cells. Large cells with their abundant cytoplasm, occupying the greater part of the cell and filled with mycobacteria leprae bacilli, highlighted by modified Fite Faraco (MFF stain, x1000)

- 7. Collagen vascular disorders: LE cells, Tart cells
 - 8. Drug induced: Podophyllin cells
 - 9. Keratinization disorders: Mantle cells, Half and half cells
 - 10. Disorders of the panniculus: Bean bag cells
 - 11. Disorders of photoimmunology: Sunburn cells
 - 12. Benign pigmented lesions: Balloon cells
 - 13. Papulosquamous disorders: Pekin cells
- B. Infectious disorders**
- 1. Bacterial: Clue cells, Downey cells, Greenblatt and Pund cells, Lepra cells, Mikulicz cells, Von Hansemann cells, Langhan giant cells
 - 2. Protozoal: Wright cells
 - 3. Viral: Koilocytes, Warthin Finkeldey cells, Mitosoid cells

REFERENCES

1. Haake AR, Holbrook K. The structure and development of skin. In: Freedberg IM, Eisen AZ, Wolff K, Austen KF, Goldsmith LA, Katz S, Fitzpatrick TB, editors. Fitzpatrick's Dermatology In General medicine. 5th ed. New York: McGraw-Hill; 2003. p. 1-109.
2. Grevelink SV, Mulliken JB. Vascular anomalies. In: Freedberg IM, Eisen AZ, Wolff K, Austen KF, Goldsmith LA, Katz S, Fitzpatrick TB, editors. Fitzpatrick's Dermatology In General medicine. 5th ed. New York: McGraw-Hill; 2003. p. 1-35.
3. Braverman IM. Ultrastructure and organization of the cutaneous microvasculature in normal and pathologic states. J Invest Dermatol 1989;93:2S-9S.
4. Braverman IM. The dermal microvascular unit: Relationship to immunological processes and dermal dendrocytes. In: Nickoloff BJ, editor. Dermal Immune System. Boca Raton: CRC Press; 1993. p. 91.
5. Bingham CO, Austen KF. The molecular and cellular biology of the mast cell. In: Freedberg IM, Eisen AZ, Wolff K, Austen KF, Goldsmith LA, Katz S, Fitzpatrick TB, editors. Fitzpatrick's

- Dermatology In General medicine. 5th ed. New York: McGraw-Hill; 2003. p. 1-33.
6. Tang CK, Tokar C. Trabecular carcinoma of the skin. *Cancer* 1978;42:2311.
 7. Gupta LK, Singhi MK. Tzanck smear: A useful diagnostic tool. *Indian J Dermatol Venereol Leprol* 2005;71:295-9.
 8. Pariser RJ. Diagnosis of spongiotic vesicular dermatitis by Tzanck smear: The "tadpole cell". *J Am Acad Dermatol* 1983;8:519-22.
 9. Durdu M, Baba M, Seçkin D. The value of Tzanck smear test in diagnosis of erosive, vesicular, bullous, and pustular skin lesions. *J Am Acad Dermatol* 2008;59:958.
 10. Lagunoff D, Ross R, Benditt EP. Histochemical and electron microscopic study in a case of Hurler's disease. *Am J Pathol* 1962;41:273-86.
 11. Hammerschmidt, Dale E, Parkin, Janet L, Brunning, Richard. Gaucher cell. *J Lab Clin Med* 1996;4:410.
 12. Naito M. Macrophage differentiation and function in health and disease. *Pathol Int* 2008;58:143-55.
 13. Braun-Falco O, Eckert F. Macroscopic and microscopic structure of xanthomatous eruptions. *Curr Probl Dermatol* 1991;20:54.
 14. Klein W, Chan E, Seykora J. Tumors of the epidermal appendages. In: Elder D, Elenitsas R, Johnson B, Murphy G, editors. *Lever's Histopathology of the skin*. 9th ed. Philadelphia: JB Lippincott; 2005. p. 874.
 15. Rivet J, Rogez C, Wechsler J. Trichoepithelioma with "monster" stromal cells. *J Cutan Pathol* 2001;28:379-82.
 16. Klein W, Chan E, Seykora J. Tumors of the epidermal appendages. In: Elder D, Elenitsas R, Johnson B, Murphy G, editors. *Lever's Histopathology of the skin*. 9th ed. Philadelphia: JB Lippincott; 2005. p. 879-80.
 17. Jaffes ES. Post-thymic T cell lymphomas. In: Jaffes ES, editors. *Surgical pathology of the lymphnodes and related organs*. 2nd ed. Philadelphia: WB Saunders; 1995. p. 360.
 18. Das P, Iyer VK, Mathur SR, Ray R. Anaplastic large cell lymphoma: A critical evaluation of cytomorphological features in seven cases. *Cytopathology* 2010;21:251-8.
 19. Dahmouh L, Hijazi Y, Barnes E, Stetler-Stevenson M, Abati A. Adult T-cell leukemia/lymphoma: A cytopathologic, immunocytochemical, and flow cytometric study. *Cancer* 2002;25:110-6.
 20. Santos JB, Farré L, Batista ES, Santos HH, Vieira MD, Bittencourt AL. The importance of flower cells for the early diagnosis of acute adult T-cell leukemia/lymphoma with skin involvement. *Acta Oncol* 2010;49:265-7.
 21. Tsukasaki K, Hermine O, Bazarbachi A, Ratner L, Ramos JC, Harrington W, et al. Definition, prognostic factors, treatment, and response criteria of adult T-cell leukemia-lymphoma: A proposal from an international consensus meeting. *J Clin Oncol* 2009;27:453-9.
 22. Stock D, McKee P, Donley B, Lakin R, Goldblum J, Howard M. Granular cell tumor of the toe: A case report. *J Foot Ankle Surg* 2009;48:358-61.
 23. Scarisbrick J, Whittacker S. Cutaneous lymphoma. In: Marcus R, John W, Sweetenham, Williams ME, editors. *Lymphoma: Pathology, diagnosis and treatment*. Cambridge: Cambridge University Press; 2007. p. 235.
 24. Ragsdale B. Tumors with fatty, muscular, osseous and cartilagenous differentiation. In: Elder D, Elenitsas R, Johnson B, Murphy G, editors. *Lever's Histopathology of the skin*. 9th ed. Philadelphia: JB Lippincott; 2005. p. 1065.
 25. Matanza-Rodriguez MI, Alvarez-Cañas MC, Gomez-Ortega JM, Fernandez F, Blanco C, Garijo F, et al. Hibernoma: A new case in the submandibular region. *Ann Saudi Med* 1996;16:670-3.
 26. Kirkham N. Tumors and cysts of the epidermis. In: Elder D, Elenitsas R, Johnson B, Murphy G, editors. *Lever's Histopathology of the skin*. 9th ed. Philadelphia: JB Lippincott; 2005. p. 853.
 27. Vonderheid EC, Bernengo MG, Burg G, Duvic M, Heald P, Laroche L, et al. Update on erythrodermic cutaneous T cell lymphoma: Report of the international Society for cutaneous lymphoma. *J Am Acad Dermatol* 2002;46:95-106.
 28. Murphy G, Schwarting R. Cutaneous Lymphomas and leukemias. In: Elder D, Elenitsas R, Johnson B, Murphy G, editors. *Lever's Histopathology of the skin*. 9th ed. Philadelphia: JB Lippincott; 2005. p. 958.
 29. Ragsdale B. Tumors with fatty, muscular, osseous and cartilagenous differentiation. In: Elder D, Elenitsas R, Johnson B, Murphy G, editors. *Lever's Histopathology of the skin*. 9th ed. Philadelphia: JB Lippincott; 2005. p. 1075.
 30. Verdolini R, Goteri G, Brancorsini D, Collina G, Simonetti O, Offidani A. Adult rhabdomyoma: Report of two cases of rhabdomyoma of the lip and of the eyelid. *Am J Dermatopathol* 2000;22:264-7.
 31. Veziroglu F, Uçkan S, Sengüven B. Adult type rhabdomyoma in a child. *Oral Oncol* 2006;42:213-6.
 32. Agamanolis DP, Dasu S, Krill CE Jr. Tumors of skeletal muscle. *Hum Pathol* 1986;17:778.
 33. Stein H. Hodgkins lymphoma: Introduction. In: Jaffe ES, Harris NL, Stein H, Vardiman JW, editors. *World Health Organization Classification of Tumours series. Pathology and Genetics of Tumours of Haematopoietic and Lymphoid Tissues*: Lyon, France: IARC Press; 2001. p. 239.
 34. Hayes TG, Rabin VR, Rosen T, Zubler MA. Hodgkin's disease presenting in the skin: Case report and review of the literature. *J Am Acad Dermatol* 1990;22:944-7.
 35. Willman JH, Golitz LE, Fitzpatrick JE. Vulvar clear cells of Tokar: Precursors of extramammary Paget's disease. *Am J Dermatopathol* 2005;27:185-8.
 36. Tokar C. Clear cells of the nipple epidermis. *Cancer* 1970;25:601-10.
 37. Shaktawat SS, Golka D. Floret-like multinucleated giant cells in neurofibroma. *Diagn Pathol* 2007;2:47.
 38. Aterman K, Remmele W, Smith M. Karl Touton and his "xanthelasmatic giant cell": A selective review of multinucleated giant cells. *Am J Dermatopathol* 1988;10:257-69.
 39. Guillou L, Calonge E, Speight P, Rosai J, Fletcher CDM. Hobnail hemangioma: A pseudomalignant vascular lesion with a reappraisal of targetoid hemosiderotic hemangioma. *Am J Surg Pathol* 1999;23:97-105.
 40. Hepburn AL. The LE cell. *Rheumatology* 2001;40:826-7.
 41. Tan EM. The LE cell and its legacy. *Clin Exp Rheumatol* 1998;16:652-8.
 42. Hargraves MM, Richmond H, Morton R. Presentation of two bone marrow elements; the "tart" cell and "LE" cell. *Proc Staff Meet Mayo Clin* 1948;23:25-8.
 43. Fisher A. Severe systemic and local reactions to topical podophyllin resin. *Cutis* 1981;28:233-6.
 44. Eady RA, Cowen T. Half-and-half cells in lichen planus. A possible clue to the origin and early formation of colloid bodies. *Br J Dermatol* 1978;98:417-23.
 45. Winkelmann RK, Bowie EJ. Hemorrhagic diathesis associated with benign histiocytic cytophagic panniculitis and systemic histiocytosis. *Arch Intern Med* 1980;140:1460-3.
 46. Sheehan JM, Young AR. The sunburn cell revisited: An update on mechanistic aspects. *Photochem Photobiol Sci* 2002;1:365-77.
 47. Mardi K, Sharma J. Metastatic balloon cell melanoma: A rare differential in the diagnosis of clear cell tumors. *Internet J Pathol* 2009;8:16.
 48. Bachner FL, Ng B, Sudilovsky D. Metastatic balloon cell melanoma: A case report. *Acta Cytol* 2005;49:543-8.
 49. Pekin TJ, Malinin TI, Zvaifler NJ. Unusual synovial fluid findings in Reiter's syndrome. *Ann Intern Med* 1967;66:677-84.
 50. Sachdeva S. Clue cell. *Indian J Dermatol Venereol Leprol* 2006;72:392-3.
 51. Downey H, McKinlay CA. Acute lymphadenosis compared with acute lymphatic leukemia. *Arch Intern Med* 1923;32:82-112.
 52. Auwaerter PG. Infectious mononucleosis in middle age. *JAMA* 1999;5:454-9.
 53. Edwin O, Wheeler, John D, Turner, Scannell J. Fever, splenomegaly and atypical lymphocytes: A syndrome observed

- after cardiac surgery utilizing a pump oxygenator. *N Engl J Med* 1962;266:454-6.
54. Pund ER, Greenblatt RB. Specific histology of granuloma inguinale. *Arch Pathol Lab Med* 1937;23:224-9.
 55. Hart G. Donovanosis. *Clin Infect Dis* 1997;25:24-32.
 56. Oliver J. Origin of the lepra cells. *J Exp Med* 1926;43:233-9.
 57. Lucas S. Bacterial diseases. In: Elder D, Elenitsas R, Johnson B, Murphy G, editors. *Lever's Histopathology of the skin*. 9th ed. Philadelphia: JB Lippincott; 2005. p. 572.
 58. Hoffman E, Loose LD, Harkin JC. The mickulicz cell in rhinoscleroma. *Am J Pathol* 1973;73:47.
 59. An T, Ferenczy A, Wilens SL, Melicow M. Observations on the formation of Michaelis-Gutmann bodies. *Hum Pathol* 1974;5:753-8.
 60. Pritchard J, Foley P, Wong H. "Langerhans and Langhans: What's misleading in a name?" *Lancet* 2003;362:922.
 61. Ramírez JR, Agudelo S, Muskus C, Alzate JF, Berberich C, Barker D, *et al.* Diagnosis of cutaneous leishmaniasis in Colombia: the sampling site within lesions influences the sensitivity of parasitologic diagnosis. *J Clin Microbiol* 2000;38:3768-73.
 62. Licona H, Bonthius DJ. Measles. In: Biller J, editor. *The Interface of Neurology and Internal Medicine*. Kluwer, Lippincott; 2007. p. 879.
 63. Xu X, Erickson L, Elder D. Diseases caused by viruses. In: Elder D, Elenitsas R, Johnson B, Murphy G, editors. *Lever's Histopathology of the skin*. 9th ed. Philadelphia: JB Lippincott; 2005. p. 664.
 64. Carlos R, Sedano HO. Multifocal papilloma virus epithelial hyperplasia. *Oral Surg Oral Med Oral Pathol* 1994;77:631-5.