

## Primary cutaneous botryomycosis

A 75-year-old woman presented with a 2-year history of progressive swelling of her right fifth toe. Physical examination showed multiple tender, smooth, round-to-oval, erythematous nodules embedded in the swollen toe without discharge [Figure 1]. Non-polarised dermoscopic examination at 10× magnification revealed scattered yellowish granules of variable size [Figure 2]. Skin biopsy taken from the nodules showed dermal deposition of granules of basophilic bacterial aggregates rimmed by an eosinophilic matrix. Gram staining revealed clustered non-filamentous Gram-positive cocci, subsequently identified as oxacillin-resistant

*Staphylococcus haemolyticus* through bacterial culture. There was no evidence of any systemic involvement. A diagnosis of primary cutaneous botryomycosis was made. The mainstay of treatment for botryomycosis consists of extended antibiotic therapy guided by sensitivity tests, frequently complemented by surgical debridement for refractory or deeply infiltrating lesions, typically resulting in favourable outcomes.<sup>1,2</sup> The patient received combination therapy involving a 3-week course of intravenous vancomycin and surgical debridement for refractory lesions, leading to the complete resolution of the lesions.




**Figure 1:** Multiple tender, smooth, round-to-oval, erythematous nodules were embedded in the swollen toe without discharge.

**How to cite this article:** Wang WY, Cheng ST. Primary cutaneous botryomycosis. Indian J Dermatol Venereol Leprol. doi: 10.25259/IJDVL\_1036\_2023

**Received:** October, 2023 **Accepted:** October, 2023 **Epub Ahead of Print:** January, 2024

**DOI:** 10.25259/IJDVL\_1036\_2023

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.



**Figure 2:** Non-polarised dermoscopic examination at 10× magnification of the nodular lesions revealed scattered yellowish granules of variable size, known as 'grains'.

#### Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent.

#### Financial support and sponsorship

Nil.

#### Conflicts of interest

There are no conflicts of interest.

#### Use of artificial intelligence (AI)-assisted technology for manuscript preparation

The authors confirm that there was no use of artificial intelligence (AI)-assisted technology for assisting in the writing or editing of the manuscript and no images were manipulated using AI.

#### *Wei-Yao Wang, Shih-Tsung Cheng<sup>1</sup>*

Department of Dermatology, Kaohsiung Medical University Hospital, Kaohsiung Medical University, Kaohsiung City, <sup>1</sup>Department of Dermatology, College of Medicine, Kaohsiung Medical University, Kaohsiung City, Taiwan.

#### Corresponding author:

Dr. Shih-Tsung Cheng,  
Department of Dermatology, College of Medicine, Kaohsiung Medical University, Kaohsiung City, Taiwan.  
tensichong@gmail.com

#### References

1. Pradhan S, Sirka CS, Panda M, Baisakh M. Cutaneous botryomycosis treated successfully with injectable ceftriaxone sodium in an immunocompetent child. *Indian J Dermatol Venereol Leprol* 2018;84:485–7.
2. Sirka CS, Dash G, Pradhan S, Naik S, Rout AN, Sahu K. Cutaneous botryomycosis in immunocompetent patients: A case series. *Indian Dermatol Online J* 2019;10:311–5.