

Striate palmoplantar keratoderma of Brünauer-Fuhs-Siemens

Received: May, 2020
Accepted: October, 2020
Published:

DOI:
10.25259/IJDVL_759_20

PMID:



Figure 1: Symmetrically distributed, linear hyperkeratosis on the flexor aspect of the fingers, the area of the right palm between the 1st and 2nd fingers and the hypothenar region of the left palm

A 48-year-old man presented with symmetric, linear hyperkeratosis on the flexor aspects of the fingers and palms [Figure 1]. He presented a family history of paternal consanguinity (fourth degree) and two sisters with similar lesions. Based on the clinical characteristics and family history, the diagnosis of striated palmoplantar keratoderma was established. The patient refused to give consent for the genetic studies.

Palmoplantar keratodermas represent a group of genodermatoses characterized by epidermal thickening.^{1,2} Striate palmoplantar keratoderma is a rare, autosomal dominant subtype of the inherited palmoplantar keratodermas presenting as linear hyperkeratosis on the palms and fingers.^{1,2} Treatment options (topical keratolytics, oral retinoids and surgical debridement) have shown moderate results.²

Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

How to cite this article: Valenzuela-Ubiña S, Jiménez-Gallo D, Linares-Barrios M. Striated palmoplantar keratoderma of Brünauer-Fuhs-Siemens. Indian J Dermatol Venereol Leprol 2021;XX:1-2.

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

Sandra Valenzuela-Ubiña, David Jiménez-Gallo, Mario Linares-Barrios

Unidad de Gestión Clínica de Dermatología Médico-Quirúrgica y Venereología, Puerta Del Mar University Hospital, Cádiz, Spain

Corresponding author:

Dr. Sandra Valenzuela-Ubiña,
Av. Ana de Viya 21, 11009, Cádiz, Spain.
sandra.vzla@hotmail.es

References

1. Kotcher LB, Jih MH, White KL, Kimyai-Asadi A. Striated palmoplantar keratoderma of Brunauer-Fohs-Siemens. *Int J Dermatol* 2001;40:644-5.
2. Tiwary AK, Chatterjee S, Mishra DK. Brunauer-Fuhs-Siemens palmoplantar keratoderma: A rare, striate type of focal palmoplantar keratoderma. *Indian J Paediatr Dermatol* 2016;17:246-8.