

Scabies surreptitius masquerading as perforating dermatosis

Dear Editor,

Scabies is a parasitic infestation caused by the mite *Sarcoptes scabiei var hominis*.¹ The diagnosis is confirmed by identifying the mite, ova, or excreta on microscopic examination. Classically, the patient presents with generalised excoriated, pruritic papules and pathognomonic burrows usually found in the finger-web space and genitalia.^{1,2} However, scabies rarely presents with an unusual or atypical morphology with no clue to the underlying infestation.³ ‘Scabies surreptitius’ or ‘scabies incognito’ is a term used for such atypical presentations of scabies.^{1,3} Here, we present a case of scabies that was not clinically suspected but was diagnosed on histopathology.

A 50-year-old immunocompetent man with poorly controlled diabetes was presented to the skin outpatient department with pruritic papular lesions all over the body. Examination revealed multiple, discrete, erythematous, perifollicular papules and nodules with central keratotic plugs and

excoriations over the trunk and proximal parts of upper and lower limbs. Few of these lesions had become secondarily infected. A large 4 cm sized abscess was noted over the left anterior axillary fold [Figure 1a]. Perforating lesions were also seen on the lower limbs and genitalia [Figure 1b]. Three clinical differential diagnoses were considered: perforating dermatosis, prurigo nodularis, and papulonecrotic tuberculid with abscess formation.

Skin biopsies were performed from two sites: one from the chest and the other from the abdomen. Grossly, the skin biopsy from the chest measured 0.6 x 0.5 x 0.2 cm, and the skin biopsy from the abdomen measured 0.5 x 0.4 x 0.3 cm. In both biopsies, the epidermis and dermis with hair follicles were identified without any subcutaneous adipose tissue.

On microscopy, the epidermis showed marked spongiosis, focal parakeratosis, and epidermal bullae with keratinocyte necrosis. The dermis showed perivascular and perifollicular chronic inflammatory infiltrates showing lymphocytes,



Figure 1a: Follicular papules and nodules with a central keratotic plug on the trunk and arms; note a pyogenic abscess in the left axilla.



Figure 1b: Perforating lesions over the lower abdomen and thighs.

How to cite this article: Agrawal A, Singal A, Devanda R, Arora VK. Scabies surreptitius masquerading as perforating dermatosis. Indian J Dermatol Venereol Leprol. doi: 10.25259/IJDVL_834_2023

Received: August, 2023 **Accepted:** October, 2023 **Epub Ahead of Print:** January, 2024

DOI: 10.25259/IJDVL_834_2023

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.




Figure 2a: Epidermal bullae and keratinocyte necrosis (Haematoxylin and eosin; 100x).




Figure 2b: Mite identified in a subcorneal burrow on serial sections. Note perivascular and perifollicular chronic inflammatory infiltrate in the dermis (Haematoxylin and eosin; 100x).

plasma cells, and occasional neutrophils with no evidence of transepithelial elimination [Figure 2a]. Findings were non-specific and no conclusive diagnosis could be made. Serial sections were advised. Incidentally, superficial corneal burrows were identified as having inflammatory exudate. One of the burrows showed the exoskeleton of a mite, *Sarcoptes scabiei* [Figure 2b] and thus a final diagnosis of scabies was made. The patient received systemic antibiotics and topical anti-scabies medication (5% permethrin) twice, with a two-week interval between treatments along with oral antihistamine. The skin lesions resolved completely within four weeks.

Scabies is highly contagious, and patients exhibit characteristic burrows overlying interdigital web spaces, wrists, elbows, lower abdomen, axilla, and breasts. However, in our case, the patient presented with perforating papules and nodules.

The diagnosis of scabies is usually made on clinical examination. However, a definitive diagnosis is made by demonstrating the mite, its eggs, or faecal pellets in the skin scrapings. A non-invasive, quick office tool, 'dermoscopy', in most cases shows a characteristic jet-plane or hang-glider appearance. A skin biopsy is rarely required. Polymerase chain reaction (PCR) and burrow ink tests are other diagnostic modalities.⁴ In our case, skin scrapings were not taken as scabies was not suspected clinically owing to its atypical presentation. The skin biopsy, which was performed to establish the diagnosis of perforating dermatoses in an

uncontrolled diabetic patient, incidentally, revealed the presence of the mite in serial sections on histopathological examination. Many different variants of scabies have been described, for example, nodular scabies, crusted scabies, and bullous scabies, among others.^{1,3} Co-existing systemic disorders or infections alter the clinical presentation of scabies.¹ Surreptitious scabies, also known as scabies incognito, refers to atypical presentations of scabies that mimic other pruritic dermatoses.^{1,2,3,5} Unusual manifestations, such as those resembling drug eruptions, Gottron papules, pityriasis rosea, and pruritic bruises associated with child abuse, have been documented in the literature, posing challenges in clinical diagnosis. Table 1 provides a summary of such surreptitious scabies cases.^{2,3,6,7}

In our patient, three clinical differential diagnoses were considered- perforating dermatoses, prurigo nodularis, and papulonecrotic tuberculid. Their clinical and histological features have been summarised in Table 2.⁵

In conclusion, the nomenclature "scabies surrepticius" aptly captures its hidden or atypical clinical manifestations, often resembling diverse dermatoses. The incidental diagnosis during the investigation of other medical conditions, as illustrated in the current case, underscores the need for a vigilant approach. Thus, a heightened index of suspicion and comprehensive tissue examinations are imperative to ensure accurate and timely diagnosis, preventing the oversight of this elusive presentation of scabies

Table 1: Some cases of Scabies Surreptitius reported in the literature

S. No	Case Details	Complaints	Lesion Morphology	Clinical Diagnosis	Confirmatory tests	Final Diagnosis
Cohen <i>et al.</i> ¹	76y/M	Recurrent itchy lesions over chest and abdomen	Red, bullous, pruritic blisters and plaques, Head and neck spared. No burrows	Drug eruption due to linezolid	HP: CICI with Eo and subepidermal vesicle with Eo. SS: Mite.	Bullous Scabies
Yoshinaga <i>et al.</i> ²	15y/F	Itchy lesions on upper extremities	Dusky red papules on proximal interphalangeal joints	Gotttron papules of dermatomyositis	SS: Mite and eggs	Scabies
Werbel <i>et al.</i> ³	91y/M	Itchy rashes on the body above the waist	Pruritic, excoriated, and erythematous nodules	Acute Folliculitis	HP: Dermal CICI with Neu. Exoskeleton of mite identified in corneal burrow.	Incognito and Nodular Scabies
Stiff <i>et al.</i> ⁶	59y/M	Lesions over the back, buttock, flanks, abdomen, face, forearms, and legs	Scaly papules and annular plaques	Pityriasis rosea	SS: Mites, ova, scybala	Scabies
Gunaric <i>et al.</i> ⁷	6y/M	Itchy bruise like lesions since 10 days	Pruritic bruises	Child Abuse	SS: Mite	Scabies
Present case	50y/M	Raised itchy lesion over the body	Pruritic raised lesions with crusting over the body. No burrows were seen [Figure 1]	Papulonecrotic tuberculid with Scrofuloderma Prurigo nodularis Perforating dermatoses	HP: Dermal CICI with Neu. Keratinocyte necrosis in subcorneal burrows along with mites [Figures 2a and 2b]	Scabies

HP: Histopathology, SS: Skin Scrapings, CICI: Chronic inflammatory cell infiltrate, Eo: Eosinophils, Neu: Neutrophils

Table 2: Clinical Differential Diagnoses

S.No	Lesion	Morphology	Histopathology
1.	Perforating Dermatoses ⁵	Perforating lesions exhibiting transepidermal elimination of substances, e.g., collagen	Four entities with distinct features: <ul style="list-style-type: none">• Acquired perforating dermatosis• Perforating folliculitis• Elastosis perforans serpiginosa• Reactive perforating collagenosis
3.	Prurigo Nodularis ⁵	Pruritic, 0.5–3 cm, hyperpigmented, firm nodules having hyperkeratotic or excoriated surface over trunk and extensor surfaces of limbs.	Ortho-hyperkeratosis with focal parakeratosis, hypergranulosis, irregular acanthosis, papillomatosis, and Irregular downward proliferation of the epidermis. Vertically oriented collagen bundles with increased numbers of fibroblasts and capillaries and perivascular chronic inflammatory cells
2.	Papulonecrotic tuberculid ⁵	Dusky papules and nodules with central necrosis. They leave varioliform scars.	Ulceration, V-shaped areas of necrosis, variable dermal thickness. Palisading histiocytes and chronic inflammatory cells and occasional granulomas. Vasculitis and fibrinoid necrosis in surrounding vessels and follicular necrosis.

Ethical approval
Institutional Review Board approval is not required.

Declaration of patient consent
The authors certify that they have obtained all appropriate patient consent.

Financial support and sponsorship
Nil.

Conflicts of interest
There are no conflicts of interest.

Use of artificial intelligence (AI)-assisted technology for manuscript preparation
The authors confirm that there was no use of artificial intelligence (AI)-assisted technology for assisting in the

writing or editing of the manuscript, and no images were manipulated using AI.

**Akanksha Agrawal, Archana Singal¹,
Rajendra Devanda¹, Vinod Kumar Arora**

Department of Pathology, ¹Department of Dermatology and STD,
University College of Medical Sciences and GTB Hospital,
New Delhi, India

Corresponding author:
Dr. Vinod Kumar Arora,
Department of Pathology, University College of Medical Sciences
and GTB Hospital, Delhi, India.
drvinodkumararora@gmail.com

References

1. Cohen PR. Scabies masquerading as bullous pemphigoid: Scabies surrepticius. *Clin Cosmet Investig Dermatol* 2017;10:317–24.
2. Yoshinaga E, Oiso N, Kawara S, Kawada A. An adolescent patient with scabies mimicking Gottron papules. *Case Rep Dermatol* 2009;2:8–12.
3. Werbel T, Hinds BR, Cohen PR. Scabies presenting as cutaneous nodules or malar erythema: Reports of patients with scabies surrepticius masquerading as prurigo nodularis or systemic lupus erythematosus. *Dermatol Online J* 2018;24:8
4. Anderson KL, Strowd LC. Epidemiology, diagnosis, and treatment of scabies in a dermatology office. *J Am Board Fam Med* 2017;30:78–84.
5. Sawant S, Gaikwad N, Hajirnis K, Vasani R. Acquired perforating collagenosis: A clinico-pathological study of ten. *Indian J Pathol Oncol* 2019;6:677–81
6. Stiff KM, Cohen PR. Scabies surrepticius: Scabies masquerading as pityriasis rosea. *Cureus* 2017;9:e1961.
7. Gunaric A, Jurisic K, Simic D, Penavic JZ, Jozic S, Goluza I. Scabies mimicking child abuse – A case report. *Psychiatr Danub* 2017;29:143–7.